



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore \triangle ABD = \triangle BCF + \triangle ACE.$$

Q. E. D.

That the three lines  $AF$ ,  $BE$ , and  $CD$  intersect in one point may be proved as follows :

Since  $EOB$  cuts the sides of  $\triangle ACM$  in the points  $N$ ,  $O$ ,  $B$ ,  $CN \times OM \times AB = AN \times CO \times BM \dots (1)$ .

Since  $AOF$  cuts the sides of  $\triangle BCM$  in  $O$ ,  $P$ ,  $F$ ,  $CO \times BP \times AM = OM \times CP \times AP \dots (2)$ .

Multiplying (1) by (2) and canceling,  $CN \times BP \times AM = AN \times BM \times CP$ .

Also demonstrated by *G. B. M. ZERR*, and the *PROPOSER*.

142. Proposed by *WILLIAM HOOVER*, A.M., Ph.D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Show that an infinite number of triangles can be inscribed in  $x^2/a^2 + y^2/b^2 - 1 = 0$  whose sides touch  $a^2x^2 + b^2y^2 = \frac{a^4b^4}{(a^2 + b^2)^2}$ .

I. Solution by the *PROPOSER*.

The curves are  $b^2x^2 + a^2y^2 - a^2b^2 = 0 \dots (1)$ ,

$$\text{and } a^2(a^2 + b^2)^2x^2 + b^2(a^2 + b^2)^2y^2 - a^4b^4 = 0 \dots (2).$$

The invariants of (1) are  $\Delta = -a^4b^4$ ,

$$\theta' = -a^2b^2(a^4 + a^2b^2 + b^4)^2$$

and of (2),  $\Delta = -a^6b^6(a^2 + b^2)^4$ ,

$$\theta' = -2a^4b^4(a^2 + b^2)^2(a^4 + a^2b^2 + b^4).$$

By the usual theory, the conditions of the problem are fulfilled if  $\theta^2 = 4\Delta\theta' \dots (3)$ , which is easily seen to be the case here.

II. Solution by *GEORGE A. OSBORNE*, Professor of Mathematics, Massachusetts Institute of Technology, Boston, Mass.

The problem is a special case of the following :

An infinite number of triangles can be inscribed in the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

whose sides touch  $\frac{x^2}{a'^2} + \frac{y^2}{b'^2} = 1$ , provided  $\frac{a'}{a} + \frac{b'}{b} = 1$ .

The problem is considered in Salmon's *Conic Sections*, Art. 376, page 342, for the general form of the conic.

$$S = ax^2 + by^2 + cz^2 + 2fyz + 2gzx + 2hxy = 0.$$

$$S' = a'x^2 + b'y^2 + c'z^2 + 2f'yz + 2g'zx + 2h'xy = 0.$$

The condition that an infinite number of triangles may be inscribed in  $S$  and circumscribed about  $S'$ , is  $\Theta'^2 = 4\Delta'\Theta \dots (1)$ .

$\Delta$ ,  $\Delta'$ , are the discriminants of  $S$ ,  $S'$ .

$$\Delta = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}, \quad \Delta' = \begin{vmatrix} a' & h' & g' \\ h' & b' & f' \\ g' & f' & c' \end{vmatrix}.$$

$$\Theta = Aa' + Bb' + Cc' + 2Ff' + 2Gg' + 2Hh',$$

$$\Theta' = A'a + B'b + C'c + 2F'f + 2G'g + 2H'h,$$

where  $A, B, C$ , etc., are minors of  $\Delta$ , and  $A', B', C'$ , etc., are minors of  $\Delta'$ .

$$\text{For } \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0 \text{ and } \frac{x^2}{a'^2} + \frac{y^2}{b'^2} - 1 = 0, \text{ we find } \Delta' = -\frac{1}{a'^2 b'^2},$$

$$\Theta = -\frac{1}{a^2 b'^2} - \frac{1}{a'^2 b^2} - \frac{1}{a^2 b^2}, \quad \Theta' = -\frac{1}{a'^2 b^2} - \frac{1}{a^2 b'^2} - \frac{1}{a'^2 b'^2}.$$

Substituting in (1), we have

$$\left( \frac{a'^2}{a^2} + \frac{b'^2}{b^2} + 1 \right)^2 = 4 \left( \frac{a'^2}{a^2} + \frac{b'^2}{b^2} + \frac{a'^2 b'^2}{a^2 b^2} \right)$$

$$\text{from which } \frac{a'}{a} \pm \frac{b'}{b} = \pm 1.$$

The only real solution for ellipses is

$$\frac{a'}{a} + \frac{b'}{b} = 1.$$

In problem 142,

$$a' = \frac{ab^2}{a^2 + b^2}, \quad b' = \frac{a^2 b}{a^2 + b^2}, \text{ from which } \frac{a'}{a} + \frac{b'}{b} = 1.$$

A very excellent demonstration was received from *G. B. M. ZERR*.

---

## CALCULUS.

---

### NOTE ON CENTER OF CURVATURE.

By *GEORGE R. DEAN, A. M.*, Professor of Mathematics, University of Missouri School of Mines and Metallurgy, Rolla, Mo.

The fact, that the point of intersection of two normals which approach each other is neither at infinity nor at the foot of the normals when they become coincident, is not plain to most students. The difficulty may be overcome, in some instances, in the following manner.

Let  $(x_1, y_1), (x_2, y_2)$  be two points of the curve;  $m_1; m_2$  the slope of the tangents at these points. The equations of the normals are then